

ON THE COMPOUND STRUCTURES OF THE NEUTRINO MASS AND CHARGE

Rasulkhozha S. Sharafiddinov

**Institute of Nuclear Physics, Uzbekistan Academy of Sciences,
Tashkent, 702132 Ulugbek, Uzbekistan**

The mass and charge of a particle correspond to the most diverse form of the same regularity of the nature of this field. As a consequence, each of all possible types of charges testifies in favor of the existence of a kind of the inertial mass. Therefore, to investigate these features we have established the compound structures of mass and charge. They can explain also the availability of fundamental differences in the masses as well as in the charges of Dirac and Majorana neutrinos.

One of sharply expressed features of the interaction of Dirac ($\nu_D \neq \bar{\nu}_D$) and Majorana ($\nu_M = \bar{\nu}_M$) neutrinos with field of emission [1, 2, 3] is the connection between these phenomena and character of the structure of fermions [4, 5, 6, 7] themselves. At the same time a question about the nature, similarity and difference of masses of neutrinos of both types [8] remains thus far not finally investigated.

The nature is, according to well known considerations, created so that all the forces in it be have the unified regularity. From this point of view, becomes possible use the Newton law of gravity as the Coulomb law and vice versa. In other words, these forces correspond to the most diverse form of the same action. Exactly the same one can as the interacting objects choose the two of neutrinos. Such a procedure, however, takes place regardless of the neutrino structure of whether it is the Dirac or the Majorana fermion. In this a hard connection is said between the inertial mass of a particle and its physical nature.

Our study of elastic scattering of electrons and their neutrinos on a spinless nucleus shows [9, 10] clearly that if the neutrino is the four - component particle ($\nu = \nu_D = \nu_e$) having a Dirac mass m_{ν_D} , it must possess both normal and anomalous electric charges. According to these data, the neutrino full electric charge in the static limit has the size

$$e_\nu = -\frac{3eG_F m_\nu^2}{4\pi^2 \sqrt{2}}, \quad e = |e|. \quad (1)$$

Such a picture leading to the flip of the neutrino spin [11] and reflects the fact that the mass and charge of a particle correspond to two form of the same regularity of the nature of its structure [9, 10].

For further purposes of a given work it is desirable to remind about the electron mass. From point of view of the classical theory of electromagnetic mass [12], the availability of the eigenenergy E_0 of the electron electrostatic field implies the existence of the electric part of the electron rest mass:

$$m_e^{em} = \frac{E_0}{c^2}.$$

The assumption has even been speaked out that all the mass of the electron is equal to its electromagnetic mass. This idea was simply called a hypothesis of field mass.

From our earlier developments, we find that a particle mass is strictly multicomponent. One of them corresponds to the electric charge and can be

called a Coulomb mass. Insofar as the electrically neutral neutrino [13] is concerned, its mass does not contain the part, at which it would have as well as an electric charge.

The difference in the masses of Dirac ($\nu = \nu_D$) and Majorana ($\nu = \nu_M$) neutrinos is observed because each of all possible types of charges of the same neutrino arises as a consequence of the availability of a kind of the inertial mass. Thereby this mechanism leads to the appearance of the united rest mass m_ν^U of the neutrino equal to all the mass of a given particle. Its general structure at the account of nonweak [14, 15] and unknown properties of the neutrino has the form

$$m_\nu^U = m_\nu^E + m_\nu^W + m_\nu^S + \dots, \quad (2)$$

where m_ν^E , m_ν^W and m_ν^S denote respectively the electric, weak and strong components of mass. Such a sight to the origination of m_ν^U quality explains the presence of the united charge e_ν^U of the neutrino equal to all the charge of a given fermion which contains the electric e_ν^E , weak e_ν^W , strong e_ν^S and some other parts:

$$e_\nu^U = e_\nu^E + e_\nu^W + e_\nu^S + \dots \quad (3)$$

So, it is seen that the mass and charge of a Dirac particle include in self both electric and unelectric components. Of course, our formula (1) characterize only a Coulomb mass dependence of the electric charge:

$$m_\nu = m_{\nu_D}^E, \quad e_\nu = e_{\nu_D}^E.$$

Using this and by following the fact that the force of Newton attraction between the two neutrinos is less than the force of their Coulomb repulsion, we find that

$$m_{\nu_D}^E > 1.53 \cdot 10^{-3} \text{ eV}, \quad (4)$$

$$e_{\nu_D}^E > 1.46 \cdot 10^{-30} \text{ e}. \quad (5)$$

It is clear, however, that the finding values of (4) and (5) are incompatible with the available laboratory data [16]. At the same time this circumstance may serve as some confirmation of the availability of compound structures of mass and charge.

Thus, it follows that if neutrinos are of electrically neutral ($\nu = \nu_M$) then $m_{\nu_M}^E = 0$, $e_{\nu_M}^E = 0$, and the size of $m_{\nu_M}^U$ and $e_{\nu_M}^U$ are reduced to the form

$$m_{\nu_M}^U = m_{\nu_M}^W + m_{\nu_M}^S + \dots, \quad (6)$$

$$e_{\nu_M}^U = e_{\nu_M}^W + e_{\nu_M}^S + \quad (7)$$

Comparing their with (2) and (3) at $\nu = \nu_D$, it is easy to observe the fundamental differences in the masses as well as in the charges of neutrinos of the different nature.

Of course, the above - noted regularities of general picture of massive neutrinos extending well known hypothesis of field mass meet with many problems which require the study of the structure and property of each of existing types of charges and masses.

Acknowledgement

I would like to express my gratitude to Professor Burkhan K. Kerimov for his numerous valuable discussions and critical remarks that was useful in improving the manuscript.

References

- [1] P.B. Pal and L. Wolfenstein, *Phys. Rev. D* **25** (1982) 766
- [2] B. Kayser, *Phys. Rev. D* **26** (1982) 1662
- [3] J.F. Nieves, *Phys. Rev. D* **26** (1982) 3152
- [4] R.B. Begzhanov and R.S. Sharafiddinov, in *Proc. Int. Conf. on Nuclear Physics, Moscow, June 16-19, 1998 (St - Petersburg, 1998)*, p.353
- [5] R.B. Begzhanov and R.S. Sharafiddinov, *Mod. Phys. Lett. A* **15** (2000) 557; *Izv. Russ. Acad. Nauk Ser. Fiz.* **64** (2000) 2221
- [6] R.S. Sharafiddinov, in *Proc. Int. Conf. on Nuclear Physics, St - Petersburg, June 14-17, 2000 (St - Petersburg, 2000)*, p.121
- [7] R.S. Sharafiddinov, *Supplementary Volume of Proc. Int. Conf. Physical Interpretations of Relativity Theory - VII. London, September 15-18, 2000* in press.
- [8] Ya.B. Zel'dovich and M.Yu. Khlopov, *Uspehi Fiz. Nauk.* **135** (1981) 45
- [9] R.S. Sharafiddinov, in *Proc. Ukrain - Russian Grav. Conf. "Gravitation, Cosmology and Relativistic Astrophysics" (November 8-11, 2000, Kharkov, Ukraine)*, p.25
- [10] R.S. Sharafiddinov, *Spacetime & Substance* **1** (2000) 176
- [11] R.S. Sharafiddinov, *Dokl. Akad. Nauk Ruz. Ser. Math. Tehn. Estest.* **7** (1998) 25
- [12] E. Fermi, *Rend. Lincei* **31** (1922) 184, 306; *Phys. Zeit.* **23** (1922) 340
- [13] E. Majorana, *Nuovo Cimento* **14** (1937) 171
- [14] L.B. Okun, *Yad. Fiz.* **45** (1987) 158
- [15] R.B. Begzhanov and R.S. Sharafiddinov, in *Proc. Int. Conf. on Nuclear Physics, Moscow, June 16-19, 1998 (St - Petersburg, 1998)*, p.354
- [16] S. Davidson, B. Campbell and K.D. Bailey, *Phys. Rev. D* **43** (1991) 2314